Initial OMP development for the South Coast rock lobster resource

S.J. Johnston and D.S. Butterworth

MARAM Department of Mathematics and Applied Mathematics University of Cape Town Rondebosch 7701 Cape Town

In order to initiate OMP development for the south coast rock lobster resource, the authors propose the following. As further proposals and, if possible, some preliminary computations on this are planned for submission for further discussion at the international stock assessment workshop at the end of November, any immediate feedback from the Rock Lobster Working Group would be appreciated.

1. Assessment Model (to serve as reference case operating model)

Model 2 (the ASPM which fits to CPUE and catch-at-age data, and which allows for time-varying selectivity – see WG/09/05/SCRL8) will be used for preliminary OMP development.

2. Management Objective

The preliminary management objective to be used when tuning alternate OMPs, is for $B_{2016}^{sp} / K^{sp} = 0.40$, which serves as a customary default for many US fisheries. A possible alternate to investigate is maintenance of current spawning biomass.

Each candidate OMP (TAC-rule) will be used to set TACs for 2006-2015 (10 year period) towards this end. Note that there may be trade-offs in relation to the period over which the target abundance level is sought to be attained.

3. Statistical framework

The outputs from a Bayesian MCMC assessment will be used as the basis for projecting the resource into the future.

4. Summary statistics

The following summary statistics will be reported. The median values and the 90% probability intervals will be provided.

C_{ave}	the average catch over the 2006-2015 period
AAV	the average inter-annual catch variation over the 2006-2016 period
B_{2016}^{sp} / K^{sp}	the final spawning biomass relative to K
B_{2016}^{sp} / B_{2006}^{sp}	the final spawning biomass relative to current spawning biomass

 B_{2016}^{exp} / B_{msy} the final exploitable biomass relative to B_{msy}

In addition similar statistics will be provided for the following assessment outputs:

B_{2006}^{sp} / K^{sp}	the current spawning biomass relative to K
$B_{2006}^{ m exp}$ / $K^{ m exp}$	the exploitable biomass relative to K
$B_{2006}^{ m exp}$ / B_{msy}	the current exploitable biomass relative to B_{msy}
MSY	the maximum sustainable yield

5. Input data for the OMP

The data available for input into an OMP will be historic catch series, CPUE and catch-at-age data.

6. Generation of Future data

For the moment, future CPUE will be the only additional data used as input into an OMP. For future years (2006+) future CPUE data will be generated as follows:

 $CPUE_{v} = qB_{v}^{\exp}e^{\varepsilon_{v}} \qquad \varepsilon_{v} \sim N(0, \sigma_{CPUE}^{2})$

7. Assumptions regarding projections into the future

For the reference case (RC):

a) Future recruitment is assumed to follow the stock-recruit curve with stochastic residuals generated from $N(0, \sigma_R^2)$ [$\sigma_R = 0.4$]

b) Future selectivity functions assume $\delta_y \sim N(0, \sigma_{sel}^2)$

8. Robustness tests

To be specified later.